

# Status of the PrimEx-II Analysis on the $\pi^0$ Lifetime Measurement

Lingling Ma  
Lanzhou University/UNCW

(On behalf of PrimEx Collaboration)

# Outline

- Physics Motivation
- Experimental Setup
- Data analysis Status
- Summary

# Physics Motivation

- $\pi^0 \rightarrow \gamma\gamma$  decay width dominated by the QCD chiral anomaly is a fundamental QCD prediction
- The decay width of  $\pi^0$  predicted by chiral anomaly is exact in chiral limit (when the light quarks are massless) :

$$\Gamma(\pi^0 \rightarrow \gamma\gamma) = \frac{\alpha^2 N_c^2 m_\pi^3}{576 \pi^3 F_\pi^2} = 7.725 \text{ eV}$$

- Corrections to the chiral anomaly prediction:

### Calculations in NLO ChPT:

- $\Gamma(\pi^0 \rightarrow \gamma\gamma) = 8.10 \text{ eV} \pm 1.0\%$

(J. Goity, et al. Phys. Rev. D66:076014, 2002)

- $\Gamma(\pi^0 \rightarrow \gamma\gamma) = 8.06 \text{ eV} \pm 1.0\%$

(B. Ananthanarayan et al. JHEP 05:052, 2002)

### Calculations in NNLO SU(2) ChPT:

- $\Gamma(\pi^0 \rightarrow \gamma\gamma) = 8.09 \text{ eV} \pm 1.3\%$


(K. Kampf et al. Phys. Rev. D79:076005, 2009)

- Calculations in QCD sum rule:

- $\Gamma(\pi^0 \rightarrow \gamma\gamma) = 7.93 \text{ eV} \pm 1.5\%$

(B.L. Ioffe, et al. Phys. Lett. B647, p. 389, 2007)

- Precision measurements of  $\Gamma(\pi^0 \rightarrow \gamma\gamma)$  at the percent level will provide a stringent test of a fundamental prediction of QCD.




# Physics Goal for PrimEx-II

## □ PrimEx-I:

$\Gamma(\pi^0 \rightarrow \gamma\gamma) = 7.82 \pm 0.14 \pm 0.17 \text{ eV}$  ( $\pm 2.8\%$  total).  
(I. Larin et al., Phys.Rev.Lett., 106:162303, 2011)


## □ PrimEx-II:

projected total uncertainty :  $\pm 1.4\%$



# PrimEx-II Experimental Setup


- ❑ Used 5.7 GeV continuous  $e^-$  beam and Hall B high resolution, high intensity photon tagging facility
- ❑ Pair spectrometer for photon flux control at high intensities.
- ❑ Two targets:  $^{12}\text{C}$  and  $^{28}\text{Si}$  (10% R.L.)
- ❑ High resolution hybrid multi-channel calorimeter (HYCAL) to detect photons from  $\pi^0$  decays
- ❑ Measured 3 absolute cross sections:
  - ✓ Primary:  $\pi^0$  production
  - ✓ Secondary: electron Compton and  $e^+e^-$  pair production



# $\pi^0$ event selection


## Measured quantities:

- ✓ initial photon energy and time
- ✓ energy and time of decayed photons
- ✓ X,Y positions of decayed photons




## Kinematical constraints:

- ✓ Conservation of energy;
- ✓ Conservation of momentum;
- ✓  $m_{\gamma\gamma}$  invariant mass




# Data Analysis Status


## Tagger timing calibration




## $\gamma\gamma$ Constraint Invariant Mass, $0.0 < \theta \leq 0.25$



## Elasticity Vs. $\gamma\gamma$ Invariant Mass, $0.0 < \theta \leq 0.25$



# Extracted $\pi^0$ Yields on $^{28}\text{Si}$ target (Preliminary Results)





# Summary

- PrimEx-II experiment was performed in Hall B to extract  $\Gamma(\pi^0 \rightarrow \gamma\gamma)$  with high precision to provide a stringent test of a fundamental QCD prediction
- Two targets:  $^{12}\text{C}$  and  $^{28}\text{Si}$
- Data analysis is in progress:
  - ✓ calibrations are done
  - ✓ Photon flux analysis is completed
  - ✓ Preliminary results on  $\pi^0$  yield and primakoff amplitude extraction on  $^{28}\text{Si}$  target are obtained
- Future plans:
  - Extract final  $\pi^0$  differential cross sections for both  $^{12}\text{C}$  and  $^{28}\text{Si}$  targets
  - Fit cross sections and extract  $\Gamma(\pi^0 \rightarrow \gamma\gamma)$
  - Finalize experimental systematic uncertainties with analyzing the electron Compton and  $e^+e^-$  pair production data

This project is supported by NSF (PHY-0079840 (MRI) ,  
PHY-1206043)and Jefferson Lab

Thank you!